

This notebook accompanies the paper: "Creative Telescoping on Multiple Sums" by Christoph Koutschan and Elaine Wong.

The paper has been submitted for publication to a special issue of the Springer-Birkhauser Journal, Mathematics in Computer Science, as part of the post-proceedings for CASC 2020.

Package Loading

The following packages are available for free download at this link: <https://www3.risc.jku.at/research/-combinat/software/ergosum/packages.html>

They should be saved in the same directory as the notebook.

```
In[ ]:= SetDirectory[NotebookDirectory[]];  
<< RISC`HolonomicFunctions`;  
<< RISC`Guess`;
```

HolonomicFunctions Package version 1.7.3 (21-Mar-2017)
written by Christoph Koutschan
Copyright Research Institute for Symbolic Computation (RISC),
Johannes Kepler University, Linz, Austria

--> Type ?HolonomicFunctions for help.

Package GeneratingFunctions version 0.8 written by Christian Mallinger
Copyright Research Institute for Symbolic Computation (RISC),
Johannes Kepler University, Linz, Austria

Guess Package version 0.52
written by Manuel Kauers
Copyright Research Institute for Symbolic Computation (RISC),
Johannes Kepler University, Linz, Austria

Definitions

```

In[ ]:= (* Original Formula (1) *)
polyG[b_, m_, s_, x_] :=
  Sum[Binomial[s, r] Binomial[k-1, r-1] (b-1) (b x)^k / (-b)^r
    Sum[(-b)^i Binomial[r-1, i], {i, 0, r-1-Max[k-m, 0]}], {k, m+s-1}, {r, s}]
(* Split Sums *)
G1[b_, m_, s_, x_] := -Sum[
  Binomial[s, r] Binomial[k-1, r-1] (b-1)^r (b x)^k / b^r, {k, m+s-1}, {r, s}]
G2[b_, m_, s_, x_] := Sum[Binomial[s, r] Binomial[k-1, r-1] (1-b) / (-b)^r
  Sum[Binomial[r-1, i] (-b)^i (b x)^k, {i, r-(k-m), r-1}],
  {k, m+1, m+s-1}, {r, s}]
(* Triple Sum No Max (8) *)
summand[b_, m_, s_, x_, k_, r_, i_] :=
  Binomial[s, r] Binomial[k-1, r-1] Binomial[r-1, i] (b-1) (b x)^k / (-b)^(r-i)
tripleG[b_, m_, s_, x_] := Sum[summand[b, m, s, x, k, r, i],
  {k, m+s-1}, {r, s}, {i, 0, r-1-(k-m)}]

```

```

In[ ]:= (* We do a few experimental sanity checks
  to confirm that our definitions are the same. *)
Timing[(Table[Together[polyG[b, m, s, x] - G1[b, m, s, x] - G2[b, m, s, x]],
  {m, 30}, {s, 30}] // Flatten // Union) === {0}]
Timing[(Table[Together[polyG[b, m, s, x] - tripleG[b, m, s, x]], {m, 30}, {s, 30}] //
  Flatten // Union) === {0}]

```

```
Out[ ]:= {137.95, True}
```

```
Out[ ]:= {135.61, True}
```

Toy Example (6)

ALERT: We prefix all functions with "te-" (for toy example) in this section to prevent repeated usage later.

```

In[ ]:= TraditionalForm[HoldForm[Sum[Binomial[n, k], {k, 5, n}]]]
teSummand[n_, k_] := Binomial[n, k];

```

```
Out[ ]//TraditionalForm=
```

$$\sum_{k=5}^n \binom{n}{k}$$

Creative Telescoping can take as input the summand OR the annihilator of the summand for the sum to be determined.

```
In[ ]:= CreativeTelescoping[Annihilator[teSummand[n, k], {S[k], S[n]}], S[k] - 1, S[n]]
```

```
Out[ ]:= {{Sn - 2}, {- $\frac{k}{-1 + k - n}$ }}
```

```
In[ ]:= {tePList, teQList} = CreativeTelescoping[teSummand[n, k], S[k] - 1, S[n]]
```

```
Out[ ]:= {{Sn - 2}, {- $\frac{k}{-1 + k - n}$ }}
```

Here, we deal with singularities and commutation.

```
In[ ]:= (* Since the sum takes on nonzero values outside of the summation
        bounds AND the operator and the summation does not commute AND
        the certificate has a singularity at one of the boundary values,
        we have a nonzero inhomogeneous part. We collect all such terms here. *)
```

```
teDeltapart[lower_, upper_] :=
```

```
  With[{delta = ApplyOreOperator[teQList[[1]], Binomial[n, k]]},
        (delta /. {k -> (upper + 1)}) - (delta /. {k -> lower})]
```

```
teCompensated[upper_] := Module[{j = n - upper},
```

```
  -Binomial[n + 1, n - j + 1] -
```

```
  Sum[(Binomial[n + 1, n + 1 - j + k] - 2 Binomial[n, n - j + k]), {k, 1, j}]
```

```
];
```

```
Table[teCompensated[i] // Together, {i, n - 3, n + 3, 1}]
```

```
Out[ ]:= { $\frac{1}{6}(-2n + 3n^2 - n^3)$ ,  $\frac{1}{2}(n - n^2)$ , -n, -1, 0, 0, 0}
```

```
In[ ]:= (* Such parts disappear if the lower limit of the sum is ≤
        0 or the upper limit is ≥ n+1, with a division by zero error for k=n. *)
```

```
(* We use the term "natural boundaries" to describe the sum
in the case that the inhomogeneous parts disappear. *)
```

```
teInhomoparts[lower_, upper_] := teCompensated[upper] + teDeltapart[lower, upper];
```

```
teInhomoparts[5, n - 1] // FullSimplify
```

```
Quiet[teInhomoparts[5, n]]
```

```
teInhomoparts[0, n + 1]
```

```
Out[ ]:= - $\frac{1}{24}(-3 + n)(-2 + n)(-1 + n)n$ 
```

```
Out[ ]:= Indeterminate
```

```
Out[ ]:= 0
```

```

In[ ]:= (* We can now find the annihilator for the inhomogeneous part and
        multiply (on the left) to P to get the annihilator for the whole sum. *)
teAnninhomo = Annihilator[teInhomoparts[5, n - 1], {S[n]}][[1]];
teAnn = teAnninhomo ** tePlist[[1]];
(* Check. *)
ApplyOreOperator[teAnn, Sum[teSummand[n, k], {k, 5, n}]] // Together

```

Out[]:= 0

We solve the recurrence to get a closed form, equivalent to Mathematica's result.

```

(* Here, we illustrate the use of Carsten Schneider's Sigma package,
which has a very nice recurrence solver. *)
Sigma`Summation`Objects`S = SigmaS;
Sigma`Summation`SumProducts`CreativeTelescoping = SigmaCT;
Quiet[Import["Sigma.m"], General::shdw]
Remove[Sigma`Summation`Objects`S,
       Sigma`Summation`SumProducts`CreativeTelescoping];

```

Sigma – A summation package by Carsten Schneider – © RISC – V 2.01 (May 12, 2016) Help

```

In[ ]:= (* Solving the recurrence and comparing closed forms. *)
teInicond = Table[Sum[teSummand[n, k], {k, 5, n}], {n, 0, 15}];
SolveRecurrence[ApplyOreOperator[teAnn, c[n]] == 0, c[n]];
teClosedall = FindLinearCombination[%, teInicond, n, 3, MinInitialValue -> 5] /.
       Sigma`Summation`Objects`Private`MyPower -> Power

```

Out[]:= $\frac{1}{24} (-24 + 3 \times 2^{3+n} - 14n - 11n^2 + 2n^3 - n^4)$

```

In[ ]:= (* The closed form is equivalent to Mathematica's version. *)
teClosedmath = Together[Simplify[Sum[Binomial[n, k], {k, 5, n}]]];
teClosedall === teClosedmath

```

Out[]:= True

Guessing

```
In[ ]:= (* We generate data necessary for guessing in all four variables. *)
(* We impose what we believe the shape of the recurrence to be,
and if such a recurrence exists, the function will find one. *)
datalist =
  (Table[polyG[b, m, s, x], {x, 2, 10}, {b, 2, 10}, {m, 10}, {s, 10}] // Together);
guessrec = GuessMultRE[datalist, {c[x, b, m, s], c[x, b, m, s + 1], c[x, b, m, s + 2],
  c[x, b, m, s + 3]}, {x, b, m, s}, 4, StartPoint -> {2, 2, 1, 1}][[1]]
```

```
Out[ ]:= (1 + m + s) (-1 + x)^2 c[x, b, m, s] -
  (-1 + x) (-4 - 2 m - 3 s + x + b x + m x + b m x + s x + b s x) c[x, b, m, 1 + s] +
  (5 + m + 3 s - 3 x - 3 b x - m x - b m x - 2 s x - 2 b s x + b x^2 + b m x^2 + b s x^2) c[x, b, m, 2 + s] +
  (2 + s) (-1 + b x) c[x, b, m, 3 + s]
```

(* Since the recurrence only depends on s,
we make some substitutions so that it looks nicer for printing. *)
(* This is the minimal order recurrence

that generates the annihilating ideal for polyG. *)
guessrec = guessrec /. (Thread[c[x, b, m, s + #] -> c[s + #]] & /@ (Range[4] - 1))

```
Out[ ]:= (1 + m + s) (-1 + x)^2 c[s] - (-1 + x) (-4 - 2 m - 3 s + x + b x + m x + b m x + s x + b s x) c[1 + s] +
  (5 + m + 3 s - 3 x - 3 b x - m x - b m x - 2 s x - 2 b s x + b x^2 + b m x^2 + b s x^2) c[2 + s] +
  (2 + s) (-1 + b x) c[3 + s]
```

Strategy Implementations for our Problem

The following implementations do not depend on each other and each section can be run on its own.
All relevant definitions for that particular section are self-contained.

Treatment of Inhomogeneous Parts: Singularities and Commutation

```
In[ ]:= (* Summand of G1 *)
f[b_, s_, x_, k_, r_] := Binomial[s, r] Binomial[k - 1, r - 1] (b - 1)^r (b x)^k / b^r;
(* Innersum of G1 *)
F1[b_, s_, x_, k_] := Sum[f[b, s, x, k, r], {r, 1, s}];
(* Creative Telescoping: Observe that we have a singularity at r =
s + 1 in the first certificate. *)
{Pinner1, Qinner1} = CreativeTelescoping[f[b, s, x, k, r], S[r] - 1, {S[s], S[k]}]
Out[ ]:= {{b s x S_s + (-1 - k) S_k + (k x - s x), (2 + k) S_k^2 + (-x - b x - k x - b k x + s x - b s x) S_k + b k x^2},
  {- (b (-k r + k r^2 + r s - r^2 s) x) / ((1 + k - r) (-1 + r - s)), b^2 k (-r + r^2) x^2 / (2 + 3 k + k^2 - 3 r - 2 k r + r^2)}}
```

```

In[ ]:= deltapart[lower_, upper_, head_, pos_] :=
  With[{delta = ApplyOreOperator[head[[pos]], f[b, s, x, k, r]],
        (delta /. {r -> (upper + 1)}) - (delta /. {r -> lower})]}
  (* We determine the delta part for the sum up to r=s-1 (i.e.,
     up to the point that we can avoid the singularity in our evaluation. *)
  deltapart[1, s - 1, Qinner1, 1] // Simplify
Out[ ]:= 
$$\frac{1}{-1 - k + s} \left( \frac{-1 + b}{b} \right)^s b^{-1+s} s^{-k+s} x (bx)^k \text{Binomial}[-1 + k, -1 + s]$$


In[ ]:= (* The original sum must be ``filled
         in'' to account for the missing term at r=s. *)
  (* This gives us the following compensation term. *)
  compensatedSing =
    -ReleaseHold[ApplyOreOperator[Pinner1[[1]], Hold[f[b, s, x, k, r]] /. r -> s]
Out[ ]:= 
$$-(-1 + b)^s b^{1-s} s(1 + s) x (bx)^k \text{Binomial}[-1 + k, -1 + s] -$$


$$(-1 + b)^s b^{-s} (bx)^k (kx - sx) \text{Binomial}[-1 + k, -1 + s] -$$


$$(-1 + b)^s b^{-s} (-1 - k) (bx)^{1+k} \text{Binomial}[k, -1 + s]$$


In[ ]:= (* We also have to concern ourselves with the compensated parts from
         any and all S[s] operators. Luckily, there is only one s-shift. *)
  compensatedSshift[coeff_, upper_, shift_, ind_List] :=
    -coeff * Sum[f[b, s + shift, x, If[ind[[1]] == 1, k + i, k], If[ind[[2]] == 1, r + i, r]],
      {i, upper + 1, upper + shift}];
  (* The annihilator for the inhomogeneous parts, together with the telescoper,
     gives the annihilator for the whole sum. *)
  inhomoparts1 = compensatedSshift[b s x, s - 1, 1, {0, 1}] +
    compensatedSing + deltapart[1, s - 1, Qinner1, 1];
  ann1 = Annihilator[inhomoparts1, {S[s], S[k]}][[1]] ** Pinner1[[1]];
  (* For the second certificate, there are no singularities,
     and the boundaries are natural,
     so the delta part is 0 and we just need to treat the commutation. *)
  inhomoparts2 = compensatedSshift[k + 2, s, 2, {0, 1}] + deltapart[1, s, Qinner1, 2];
  ann2 = Annihilator[inhomoparts2, {S[s], S[k]}][[1]] ** Pinner1[[2]];
  Support[{ann1, ann2}]
Out[ ]:= 
$$\left\{ \left\{ S_s^2 S_k, S_s S_k^2, S_k^3, S_s^2, S_s S_k, S_k^2, S_s, S_k, 1 \right\}, \left\{ S_k^3, S_k^2, S_k, 1 \right\} \right\}$$


In[ ]:= (* We now confirm that our annihilators
         are correct and indeed annihilates our sum. *)
  FullSimplify[ApplyOreOperator[{ann1, ann2}, Sum[f[b, s, x, k, r], {r, 1, s}]]]
Out[ ]:= {0, 0}

```

```

In[ ]:= (* We confirm that our computations are correct. *)
OreReduce[{ann1, ann2}, Pinner1]

Out[ ]:= {0, 0}

In[ ]:= (* In fact, the telescopers are the correct
annihilators already due to a natural boundaries argument. *)
Annihilator[F1[b, s, x, k], {S[s], S[k]}] === Pinner1

Out[ ]:= True

```

Treatment of Inhomogeneous Parts: Closure

```

In[ ]:= (* G1 *)
G1[b_, m_, s_, x_] :=
  -Sum[Binomial[s, r] Binomial[k - 1, r - 1] (b - 1)^r (b x)^k / b^r, {k, m + s - 1}, {r, s}]
(* Subs for testing. *)
subs = {b -> 31, m -> 15, x -> 5 / 7};
checksubs = {31, 15, s, x};
(* Summand of G1 *)
f[b_, s_, x_, k_, r_] := Binomial[s, r] Binomial[k - 1, r - 1] (b - 1)^r (b x)^k / b^r;
(* Innersum of G1 *)
F1[b_, s_, x_, k_] := Sum[f[b, s, x, k, r], {r, 1, s}];
(* Creative Telescoping *)
{Pinner1, Qinner1} = CreativeTelescoping[f[b, s, x, k, r], S[r] - 1, {S[s], S[k]}];
{Pouter1, Qouter1} = CreativeTelescoping[Pinner1, S[k] - 1, {S[s]}]

Out[ ]:= {{(1 - b x) S_s + (-1 + x)}, {1 + k / (b s x) S_k + (-k + s - b s) / (b s)}}

```

Symbolic Computation : Here we deal with unnatural boundaries and the fact that the telescoper does not commute with the sum and make the appropriate adjustments

```

In[ ]:= (* We first apply the certificates and then
determine the collapsing sum (up to k=m+s-1). *)
outercert1 = With[{delta = ApplyOreOperator[Qouter1[[1]], F1[b, s, x, k]]},
(delta /. k -> (s+m)) - (delta /. k -> 1)];
(* Next, we determine the annihilator of the collapsed sum and
extra stuff that was added to Sum[P*F[s,k],{k,1,m+s-1}]
so that what remains is P*Sum[F[s,k],{k,1,m+s-1}]. *)
outerstuff1 = (1 - b x) F1[b, s + 1, x, m + s];
(* The inhomogeneous part is the combination
of stuff we want to remove and the telescoped sum. *)
inhom1 = FunctionExpand[-outerstuff1 + outercert1]

```

$$\text{Out[]} = -\frac{(-1+b)(-1+s-bs)x}{b} - \frac{(-1+b)(1+b-s+bs)x}{b} + \frac{1}{b^2 x}$$

$$\begin{aligned}
& (-1+b)(1+m+s)(bx)^{1+m+s} \text{Hypergeometric2F1}\left[1-s, -m-s, 2, \frac{-1+b}{b}\right] + \frac{1}{b^2} \\
& (-1+b)(-m-bs)(bx)^{m+s} \text{Hypergeometric2F1}\left[1-s, 1-m-s, 2, \frac{-1+b}{b}\right] - \frac{1}{b} \\
& (-1+b)(1+s)(bx)^{m+s}(1-bx) \text{Hypergeometric2F1}\left[1-m-s, -s, 2, \frac{-1+b}{b}\right]
\end{aligned}$$

We now illustrate how inhom1 can be simplified and rewritten so that we can use the closure property of applying an operator.

```

In[ ]:= (* The first two terms do not contain 2F1s so they will be
easy to treat. We therefore separate inhom1 into two parts. *)
inhom1first = inhom1[[1 ;; 2]];
inhom1last = inhom1[[3 ;; 5]];
(* We rewrite inhom1last as a combination of some shifts (by observation). *)
hg[s_, m_] := (b x)^(m+s) * Hypergeometric2F1[1-s, 2-m-s, 2, (b-1)/b]
pf1 = HoldForm[(-1+b)(1+m+s)/(b^2 x * b x)];
pf2 = HoldForm[(-1+b)(-m-bs)/(b^2 * b x)];
pf3 = HoldForm[(-1+b)(1+s)(-1+bx)/(b * b x)];
(* This is the new operator constructed from the above observations. *)
op = pf1 S[m]^2 + pf2 S[m] + pf3 S[s];
applyop = ReleaseHold[ApplyOreOperator[op, hg[s, m]]];
(* We can see that our operator gives us the correct expression. *)
Simplify[inhom1last] === Simplify[applyop]

```

Out[]:= True

This allows us to compute the annihilator for inhom1 by computing the separate annihilators and applying the closure property.


```

In[ ]:= anninhom1first = Annihilator[inhom1first, {S[s]}];
anninhom1last =
  DFiniteOreAction[Annihilator[hg[s, m], {S[m], S[s]}], ReleaseHold[op]];
R1 = DFinitePlus[anninhom1first,
  {ToOrePolynomial[anninhom1last[[2]], OreAlgebra[S[s]]]}];
(* We now finally have a good annihilator for polyfirst. *)
annouter1 = (#** Pouter1[[1]]) & /@ R1;
(* The support gives an idea of the operators
  that we are dealing with in our annihilating ideal. *)
Support[annouter1]
Out[ ]:= {{S_s^4, S_s^3, S_s^2, S_s, 1}}

In[ ]:= (* Another quick check confirms that we are on the right track. *)
Table[ApplyOreOperator[#, (G1@@checksubs) /. x -> 5 / 7] & /@ annouter1, {s, 30}];
(% /. subs // Flatten // Union) === {0}

Out[ ]:= True

```

Speedup: DFiniteSub

```

In[ ]:= (* G2 *)
G2[b_, m_, s_, x_] :=
  Sum[Binomial[s, r] Binomial[k-1, r-1] (1-b) / (-b)^r Sum[Binomial[r-1, i]
    (-b)^i (b x)^k, {i, r-(k-m), r-1}], {k, m+1, m+s-1}, {r, s}];
subs = {b -> 31, m -> 15, x -> 5 / 7};
checksubs = {31, 15, s, x};
(* Inner Summand of G2 *)
h[s_, k_, r_] := Binomial[s, r] Binomial[k-1, r-1] (1-b) / (-b)^r
  Sum[Binomial[r-1, i] (-b)^i (b x)^k, {i, r-(k-m), r-1}];
(* Inner Sum of G2 *)
H[s_, k_] := Sum[h[s, k, r], {r, 1, s}];
(* Creative Telescoping *)
{Pinner2, Qinner2} = CreativeTelescoping[h[s, k, r], S[r]-1, {S[k], S[s]}];
(* Timing[{Pouter2, Qouter2}=CreativeTelescoping[Pinner2, S[k]-1, {S[s]}]; *)
(* This took about 70 seconds so we save
  it and then import it for use in this notebook. *)
{Pouter2, Qouter2} = Import["ctouter2.m"];

```

Symbolic Computation I: Here we deal with the fact that the telescoper does not commute with the sum and make the appropriate adjustments.

```

In[ ]:= (* We first apply the certificates and then
         determine the collapsing sum (up to k=m+s-1). *)
outercert2 = With[{delta = ApplyOreOperator[Qouter2, HH[s, k]],
                 (delta /. k -> (m + s)) - (delta /. k -> (m + 1))}],
(* Next we determine the stuff that must be removed to take into account
the extra shifts when moving the telescopers outside of the sum. *)
Pouter2coeff = OrePolynomialListCoefficients[Pouter2[[1]]];
outerstuff2 = Total[Pouter2coeff[[1 ;; 3]] *
                  Table[Sum[HH[s + i, k], {k, m + s, m + s + i - 1}], {i, 3, 1, -1}]];
(* The inhomogeneous part is a combination of stuff we want to
remove and the telescoped sums (from the delta). *)
inhom2 = -outerstuff2 + outercert2;

```

Symbolic Computation II: Here we try to avoid Mathematica's use of 2F1s and DifferenceRoots by rewriting all the inhomogeneous parts as an operator.

```

In[ ]:= (* We first note that the pieces that DO NOT have 2F1s evaluate to 0:
         this corresponds to evaluating the sum
         with the certificates at the lower boundary. *)
posfree = Position[FreeQ[#, Hypergeometric2F1] & /@
                 (List@@ inhom2[[1]] /. HH -> H), True] // Flatten;
inhom2[[1, posfree]] /. HH -> H // Cancel // FunctionExpand

```

Out[]:= 0

```

In[ ]:= (* We remove those zero pieces and proceed to
         determine an annihilator for the rest of inhom2. *)
inhom2rest = (inhom2[[1, Complement[Range[9], posfree]]]);
(* We construct an operator for these parts (this is done by observation). *)
newop = Qouter2 - Sum[Coefficient[Pouter2[[1]], S[s], i] **
                    S[s]^i ** Sum[S[k]^j, {j, 0, i - 1}], {i, 3}];
(* Let's make sure our operator is correct after the
application and substitution. *)
Together[inhom2rest - ApplyOreOperator[newop, HH[s, k]] /. k -> (m + s)] === {0}

```

Out[]:= True

Symbolic Computation III: Finally, we take control of the substitution by deciding which monomial to use first.

```

(* This is the long way and we keep
it here so that the timings can be compared. *)
(* Timing[R2act=DFiniteOreAction[Pinner2,newop];] *)
(* The above took 331 seconds. *)
(* Timing[R2=DFiniteSubstitute[R2act,{k->(m+s)}];] *)
(* This above took 107860 seconds!! *)
(* annouter2={R2[[1]]**Pouter2[[1]]}; *)

```

```

In[ ]:= (* Here, we are taking control of the substitution. *)
(* First, we compute the operators that act on our inhomogeneous parts. *)
(* Timing[R2act=DFiniteOreAction[Pinner2,newop];] *)
(* The above took 331 seconds so we save and reload. *)
R2act = Import["ctR2act.m"];
(* The following information allows us to determine
the holonomic rank and shape/size of our operator. *)
{# /@ R2act} & /@ {ByteCount, Support}
Out[ ]:= {{19199856, 9133192, 22585680}, {{Ss2, Sk, Ss, 1}, {Sk Ss, Sk, Ss, 1}, {Sk2, Sk, Ss, 1}}}

In[ ]:= (* We also identify the set of irreducible monomials
of the Groebner basis for the annihilating ideal. *)
UnderTheStaircase[R2act]
Out[ ]:= {1, Ss, Sk}

In[ ]:= (* We now do DFiniteSub by hand. *)
(* To do this we will construct an operator T with the desired properties. *)
(* In particular, it should be generated by the Groebner basis of R2act and have
support {1, S[s]S[k], (S[s]S[k])^2}. We do this using the command OreReduce. *)
(* We also apply the substitution already. *)
red00 = {0, 1};
red11 = OrePolynomialListCoefficients[
OreReduce[S[k] S[s], Together[R2act], OrePolynomialSubstitute → {k → m + s}]];
(* Speedup actually occurs here. *)
Timing[red22a = OreReduce[S[k]^2 S[s]^2, {Together[R2act[[2]]}],
OrePolynomialSubstitute → {k → m + s}]; Print["Timing for red22a"];]
red22a >> "red22a.m";
coeffs = Together[OrePolynomialListCoefficients[R2act[[3]]] /. k → m + s];
coeffs1 = Together[coeffs[{{2, 4}}] / coeffs[[1]]];
coeffs2 = Together[LeadingCoefficient[red22a] * coeffs1];
Timing[red22 = Together[Rest[OrePolynomialListCoefficients[red22a]] - coeffs2];
Print["Timing for red22"];]
red22 >> "red22.m";

Timing for red22a
Out[ ]:= {1004.67, Null}

Timing for red22
Out[ ]:= {3211.57, Null}

```

```

In[ ]:= (* Next, from our ansatz for T,
we solve the corresponding system for the coefficients. *)
sys = Transpose[{red00, red11, red22}];
Timing[R2 = NormalizeVector[NullSpace[sys][[1]]] // Reverse;
Print["Timing for solving the linear system."];]
R2 = ToOrePolynomial[R2.{S[s]^2, S[s], 1}, OreAlgebra[S[s]]];
(* The whole procedure takes about 4200 seconds so we save it and this
can be imported by removing the comments from the following code. *)
(* R2=Import["ctR2.m"]; *)
(* This now gives us the desired annihilator for G2. *)
annouter2 = {R2 ** Pouter2[[1]]};
UnderTheStaircase[annouter2]

Timing for solving the linear system.

Out[ ]:= {1.10798, Null}

Out[ ]:= {1, Ss, Ss2, Ss3, Ss4}

In[ ]:= (* We perform a quick sanity check to see
that this is indeed the correct annihilator for G2. *)
Table[ApplyOreOperator[#, (G2 @@ checksubs) /. x → 5 / 7] & /@ annouter2, {s, 30}];
((% /. subs) // Flatten // Union) === {0}

Out[ ]:= True

```

Speedup: Gamma Insertions

```

In[ ]:= (* Here is the guessed recurrence that we will use to compare our result. *)
guessrecpoly = ToOrePolynomial[
  (1+m+s) (-1+x)^2 c[s] - (-1+x) (-4-2m-3s+x+bx+mx+bm x+s x+bs x) c[1+s] +
  (5+m+3s-3x-3bx-mx-bm x-2s x-2bs x+bx^2+bm x^2+bs x^2) c[2+s] +
  (2+s) (-1+bx) c[3+s], c[s]];
(* Here is the summand for the triple sum. *)
summand[m_, s_, b_, x_, k_, r_, i_] := Binomial[s, r] * Binomial[k-1, r-1] *
  (- (1-b) ^ (1-r)) * (-b) ^ (i) * Binomial[r-1, i] * ((b-1)/b) ^ r * (bx) ^ k;
(* New summand with one or two gamma factors. *)
summandTwoGammas[m_, s_, b_, x_, k_, r_, i_] :=
  Binomial[s, r] * Binomial[k-1, r-1] * (- (1-b) ^ (1-r)) * (-b) ^ (i) *
  Binomial[r-1, i] * ((b-1)/b) ^ r * (bx) ^ k * (Gamma[k+eps]/Gamma[k])
  (Gamma[r-i-(k-m)+eps]/Gamma[r-i-(k-m)])
summandOneGamma[m_, s_, b_, x_, k_, r_, i_] := Binomial[s, r] *
  Binomial[k-1, r-1] * (- (1-b) ^ (1-r)) * (-b) ^ (i) * Binomial[r-1, i] *
  ((b-1)/b) ^ r * (bx) ^ k * (Gamma[r-i-(k-m)+eps]/Gamma[r-i-(k-m)])
(* Triple sum: this is the same as G and Gnomax. *)
T[m_, s_] := Hold[
  Sum[summand[m, s, b, x, k, r, i], {k, 1, m+s-1}, {r, 1, s}, {i, 0, r-1-(k-m)}]];
(* This is our corrected sum with Gammas on a slightly enlarged region. *)
Tnew2[m_, s_] := Hold[Sum[summandTwoGammas[m, s, b, x, k, r, i],
  {k, -1, m+s+1}, {r, -1, s+2}, {i, -2, r+1-(k-m)}]];
Tnew1[m_, s_] := Hold[Sum[summandOneGamma[m, s, b, x, k, r, i],
  {k, -1, m+s+1}, {r, -1, s+2}, {i, -2, r+1-(k-m)}]];


```

Figures

```

In[ ]:= (* The following figure gives a visual representation of how we
distinguish the points we like from the points we don't like. *)
(* Blue dots: the summand is non-
zero AND properly within the given boundaries. *)
(* Red dots: the summand is non-zero and outside of the given boundaries. *)
(* Empty: The summand is zero. *)
(* Goal: Manipulate the summand with
gamma corrections until we only see the blue dots. *)
limit = 5;
With[{s = 10, m = 15, b = 7, x = 1/2},
Graphics3D[Table[
  If[summand[m, s, b, x, k, r, i] == 0, {},
    {If[1 ≤ k ≤ m + s - 1 && 1 ≤ r ≤ s && 0 ≤ i ≤ r - 1 - (k - m), Blue, Red],
     Sphere[{k, r, i}, 0.3]}]
  , {k, -limit, m + s + limit}, {r, -limit, s + limit}, {i, -limit, s + limit}],
  Axes → True, AxesLabel → {"k", "r", "i"}
]]


```



```

In[ ]:= (* We can see that our "bad" (red) points go away after applying the gammas. *)
With[{s = 10, m = 15, b = 7, x = 1/2},
Graphics3D[Table[
  If[(summandTwoGammas[m, s, b, x, k, r, i] /. eps -> 0) === 0,
    {}, {If[1 ≤ k ≤ m + s - 1 && 1 ≤ r ≤ s && 0 ≤ i ≤ r - 1 - (k - m), Blue, Red]},
    Sphere[{k, r, i}, 0.3]}]
, {k, -limit, m + s + limit}, {r, -limit, s + limit}, {i, -limit, s + limit}],
Axes -> True, AxesLabel -> {"k", "r", "i"}
]]

```



Result and Timings for Two Gamma Insertions (+ Treatment of Singularities)

First we derive the annihilator for the triple sum by applying creative telescoping three times.

```

In[ ]:= Timing[
  cti2 = CreativeTelescoping[
    summandTwoGammas[m, s, b, x, k, r, i], S[i] - 1, {S[r], S[k], S[s]}];
  ctr2 = CreativeTelescoping[cti2[[1]], S[r] - 1];
  ctk2 = FindCreativeTelescoping[ctr2[[1]], {S[k] - 1};
]
Out[ ]:= {666.56, Null}

```

```

In[ ]:= (* The result is not too bad: a fourth-order recurrence in s. *)
Factor[ctk2[[1, 1]] /. eps -> 0]
(* And this recurrence lies in the ideal generated by the guessed one,
and this completes the proof. *)
OreReduce[ctk2[[1, 1]] /. eps -> 0, {guessrecpoly}]

```

$$\begin{aligned}
\text{Out[]} = & - (2 + s) (3 + s) (-1 + b x) S_s^4 - \\
& (2 + s) (9 + m + 4 s - 5 x - 6 b x - m x - b m x - 2 s x - 3 b s x + 2 b x^2 + b m x^2 + b s x^2) S_s^3 + \\
& (19 + 5 m + 21 s + 3 m s + 6 s^2 - 21 x - 7 b x - 7 m x - 3 b m x - 22 s x - \\
& 9 b s x - 4 m s x - 2 b m s x - 6 s^2 x - 3 b s^2 x + 4 x^2 + 5 b x^2 + 2 m x^2 + \\
& 3 b m x^2 + 4 s x^2 + 6 b s x^2 + m s x^2 + 2 b m s x^2 + s^2 x^2 + 2 b s^2 x^2) S_s^2 - \\
& (-1 + x) (-8 - 4 m - 11 s - 3 m s - 4 s^2 + 5 x + b x + 3 m x + b m x + 6 s x + 2 b s x + \\
& 2 m s x + b m s x + 2 s^2 x + b s^2 x) S_s + (1 + s) (1 + m + s) (-1 + x)^2
\end{aligned}$$

```
Out[ ]:= 0
```

Next, we certify that the recurrence is correct by showing that the inhomogeneous parts are 0 (expected since we have our natural boundaries). However, there are still singularities in the certificates so it is still a nontrivial (and not yet automated task) to certify.

```

In[ ]:= (* Here are the operators we have to worry about
in the telescopers and location of singularities. *)
FindSing[var_, ops_] :=
  {#[ops[[1]] /. eps -> 0] & /@ {Variables, Support}, Solve[# == 0, var] & /@
  (Denominator /@ (ApplyOreOperator[Flatten[ops[[2]]], c[var]] // Together))}
FindSing[i, cti2]
FindSing[r, ctr2]
FindSing[k, ctk2]

```

$$\text{Out[]} = \left\{ \left\{ \{b, k, m, r, s, x, S[k], S[r], S[s]\}, \{S_s, 1\}, \{S_r, S_k, 1\}, \{S_k^2, S_k, 1\} \right\}, \right. \\
\left. \left\{ \{i \rightarrow r\}, \{i \rightarrow -k + m + r\}, \{i \rightarrow -1 + \text{eps} - k + m + r\} \right\} \right\}$$

$$\text{Out[]} = \left\{ \left\{ \{b, k, m, s, x, S[k], S[s]\}, \{S_s^2, S_k, S_s, 1\}, \{S_k S_s, S_k, 1\}, \{S_k^2, S_k, S_s, 1\} \right\}, \right. \\
\left. \left\{ \{r \rightarrow 1 + s\}, \{r \rightarrow 2 + s\}, \{r \rightarrow 1 + s\}, \{r \rightarrow 1 + k\}, \{r \rightarrow 2 + k\}, \{r \rightarrow 1 + s\} \right\} \right\}$$

$$\text{Out[]} = \left\{ \left\{ \{b, m, s, x, S[s]\}, \{S_s^4, S_s^3, S_s^2, S_s, 1\} \right\}, \right. \\
\left. \left\{ \{k \rightarrow -\text{eps}\}, \{k \rightarrow 1 + m + s\}, \{k \rightarrow 2 + m + s\}, \{k \rightarrow 3 + m + s\} \right\} \right\}$$


```
(* Here we certify the first CT by confirming that the inhomogeneous
parts are 0. We only need to check it for the second certificate. *)
deltaparti[certs_, lower_, upper_] :=
  (With[{delta = ApplyOreOperator[certs, S2C[m, s, b, x, k, r, i]]},
    (delta /. {i -> (upper + 1)}) - (delta /. {i -> lower})] /. S2C -> summandTwoGammas
  compparts2 = ((ApplyOreOperator[cti2[[1, 2]], S2C[m, s, b, x, k, r, i]] /.
    {i -> r - 1 - (k - m)}) +
    ApplyOreOperator[cti2[[1, 2]], S2C[m, s, b, x, k, r, i]] /. {i -> r - 1 - (k - m)}][[
      2]]) /. S2C -> summandTwoGammas;
FullSimplify[(deltaparti[cti2[[2, 2]], 0, r - 2 - (k - m)] - compparts2) /. eps -> 0] === 0
```

Out[8]= True

```
In[9]:= (* Here we certify the second CT by
confirming that the inhomogeneous parts are 0. *)
deltapartr[certs_, lower_, upper_] := (With[{delta =
  ApplyOreOperator[certs, Sum[S2C[m, s, b, x, k, r, i], {i, 0, r - 1 - (k - m)}]]},
  (delta /. {r -> (upper + 1)}) - (delta /. {r -> lower})]
  comppartsr[1] = (ApplyOreOperator[ctr2[[1, 1]],
    Sum[S2C[m, s, b, x, k, r, i], {i, 0, r - 1 - (k - m)}]] /. {r -> s}) +
  (ApplyOreOperator[ctr2[[1, 1]], Sum[S2C[m, s, b, x, k, r, i],
    {i, 0, r - 1 - (k - m)}]] /. {r -> s + 1})[[3 ;; 4]] +
  (ApplyOreOperator[ctr2[[1, 1]], Sum[S2C[m, s, b, x, k, r, i],
    {i, 0, r - 1 - (k - m)}]] /. {r -> s + 2})[[4]];
  comppartsr[2] = (ApplyOreOperator[ctr2[[1, 2]],
    Sum[S2C[m, s, b, x, k, r, i], {i, 0, r - 1 - (k - m)}]] /. {r -> s}) +
  (ApplyOreOperator[ctr2[[1, 2]], Sum[S2C[m, s, b, x, k, r, i],
    {i, 0, r - 1 - (k - m)}]] /. {r -> s + 1})[[3]];
  comppartsr[3] = (ApplyOreOperator[ctr2[[1, 3]],
    Sum[S2C[m, s, b, x, k, r, i], {i, 0, r - 1 - (k - m)}]] /. {r -> s}) +
  (ApplyOreOperator[ctr2[[1, 3]], Sum[S2C[m, s, b, x, k, r, i],
    {i, 0, r - 1 - (k - m)}]] /. {r -> s + 1})[[4]];
  (Table[{Simplify[deltapartr[ctr2[[2, i]], 1, s - 1] - comppartsr[i]] /.
    {S2C -> summandTwoGammas} /. {eps -> 0}} // FunctionExpand //
    FullSimplify, {i, 3}] // Flatten // Union) === {0}
```

Out[9]= True

Result and Timings for One Gamma Insertion (+ Treatment of Singularities)

First we derive the annihilator for the triple sum by applying creative telescoping three times.

```

In[ ]:= Timing[
  cti1 = CreativeTelescoping[
    summandOneGamma[m, s, b, x, k, r, i], S[i] - 1, {S[r], S[k], S[s]};
  ctr1 = CreativeTelescoping[cti1[[1]], S[r] - 1];
  ctk1 = FindCreativeTelescoping[ctr1[[1]], {S[k] - 1};
]
Out[ ]:= {26.6068, Null}

In[ ]:= (* The result is exactly the third order recurrence that we guessed. *)
ctk1[[1, 1]] /. eps → 0
-% === guessrecpoly
Out[ ]:= (2 + s - 2 b x - b s x) S_s^3 +
(-5 - m - 3 s + 3 x + 3 b x + m x + b m x + 2 s x + 2 b s x - b x^2 - b m x^2 - b s x^2) S_s^2 +
(4 + 2 m + 3 s - 5 x - b x - 3 m x - b m x - 4 s x - b s x + x^2 + b x^2 + m x^2 + b m x^2 + s x^2 + b s x^2) S_s +
(-1 - m - s + 2 x + 2 m x + 2 s x - x^2 - m x^2 - s x^2)

Out[ ]:= True

In[ ]:= (* So of course this recurrence lies in the ideal generated by the guessed one,
and this completes the proof. *)
OreReduce[ctk1[[1, 1]] /. eps → 0, {guessrecpoly}]
Out[ ]:= 0

```

Next, we certify that the recurrence is correct by showing that the inhomogeneous parts are 0 (expected since we have our natural boundaries). However, there are still singularities in the certificates so it is still a nontrivial (and not yet automated task) to certify.

```

In[ ]:= (* Here are the operators we have to worry about
in the telescopers and location of singularities. *)
FindSing[var_, ops_] :=
  {#[ops[[1]] /. eps → 0] & /@ {Variables, Support}, Solve[#[# == 0, var] & /@
    (Denominator /@ (ApplyOreOperator[Flatten[ops[[2]]], c[var]] // Together))}]
FindSing[i, cti1]
FindSing[r, ctr1]
FindSing[k, ctk1]
Out[ ]:= {{{b, k, m, r, s, x, S[k], S[r], S[s]}, {{S_s, 1}, {S_r, S_k, 1}, {S_k^2, S_k, 1}}},
  {{}}, {{i → r}, {i → -k + m + r}}, {{i → -1 + eps - k + m + r}}}}
Out[ ]:= {{{b, k, m, s, x, S[k], S[s]}, {{S_s^2, S_k, S_s, 1}, {S_k S_s, S_k, 1}, {S_k^2, S_k, S_s, 1}}},
  {{r → 1 + s}, {r → 2 + s}}, {{r → 1 + s}}, {{r → 1 + k}, {r → 2 + k}, {r → 1 + s}}}}
Out[ ]:= {{{b, m, s, x, S[s]}, {{S_s^3, S_s^2, S_s, 1}}}, {{k → 1 + m + s}, {k → 2 + m + s}}}

```

```

In[ ]:= (* Here we certify the first CT by confirming that the inhomogeneous
parts are 0. We only need to check it for the second certificate. *)
deltaparti[certs_, lower_, upper_] :=
  (With[{delta = ApplyOreOperator[certs, S1C[m, s, b, x, k, r, i]]},
    (delta /. {i -> (upper + 1)} - (delta /. {i -> lower}})] /. S1C -> summandOneGamma
  compparts2 = ((ApplyOreOperator[cti1[[1, 2]], S1C[m, s, b, x, k, r, i]] /.
    {i -> r - 1 - (k - m)}) +
    ApplyOreOperator[cti1[[1, 2]], S1C[m, s, b, x, k, r, i]] /. {i -> r - 1 - (k - m)}][[
      2]]) /. S1C -> summandOneGamma;
FullSimplify[(deltaparti[cti1[[2, 2]], 0, r - 2 - (k - m)] - compparts2) /. eps -> 0] === 0

Out[ ]:= True

```

```

In[ ]:= (* Here we certify the second CT by
confirming that the inhomogeneous parts are 0. *)
deltapartr[certs_, lower_, upper_] := (With[{delta =
  ApplyOreOperator[certs, Sum[S1C[m, s, b, x, k, r, i], {i, 0, r - 1 - (k - m)}]]},
  (delta /. {r -> (upper + 1)} - (delta /. {r -> lower}})]
  comppartsr[1] = (ApplyOreOperator[ctr1[[1, 1]],
    Sum[S1C[m, s, b, x, k, r, i], {i, 0, r - 1 - (k - m)}]] /. {r -> s}) +
  (ApplyOreOperator[ctr1[[1, 1]], Sum[S1C[m, s, b, x, k, r, i],
    {i, 0, r - 1 - (k - m)}]] /. {r -> s + 1})[[3 ;; 4]] +
  (ApplyOreOperator[ctr1[[1, 1]], Sum[S1C[m, s, b, x, k, r, i],
    {i, 0, r - 1 - (k - m)}]] /. {r -> s + 2})[[4]];
  comppartsr[2] = (ApplyOreOperator[ctr1[[1, 2]],
    Sum[S1C[m, s, b, x, k, r, i], {i, 0, r - 1 - (k - m)}]] /. {r -> s}) +
  (ApplyOreOperator[ctr1[[1, 2]], Sum[S1C[m, s, b, x, k, r, i],
    {i, 0, r - 1 - (k - m)}]] /. {r -> s + 1})[[3]];
  comppartsr[3] = (ApplyOreOperator[ctr1[[1, 3]],
    Sum[S1C[m, s, b, x, k, r, i], {i, 0, r - 1 - (k - m)}]] /. {r -> s}) +
  (ApplyOreOperator[ctr1[[1, 3]], Sum[S1C[m, s, b, x, k, r, i],
    {i, 0, r - 1 - (k - m)}]] /. {r -> s + 1})[[4]];
  (Table[{Simplify[deltapartr[ctr1[[2, i]], 1, s - 1] - comppartsr[i]] /.
    {S1C -> summandOneGamma} /. {eps -> 0}} // FunctionExpand //
    FullSimplify, {i, 3}] // Flatten // Union) === {0}

Out[ ]:= True

```

Export to PDF

```

In[5]:= SetDirectory[NotebookDirectory[]];
Export["sums_casc2020.pdf", Get["sums_casc2020.nb"] /. ButtonBox[___] -> " Help"];

```